

Best Practice Adoption Level Calculation

Updated: April 2023

#	Best Practice	Measure	Prescribing Provider		ED Provider		Non-Prescribing Provider	
			Medium	High	Medium	High	Medium	High
Adoption level								
1	ROA	80% of ROAs complete and timely (provider and worker sections)	X	X	X	X	X	X
2	APF	80% of all claims have an APF with the ROA (first visit)	X	X	X	X	X	X
3	APF	80% of all time-loss claims have a 2 nd APF within four weeks of claim established date		X				X
4	APF	80% of all APFs received in two L&I business days		X		X		X
5	Two-Way Communication	25% of all claims have two-way communication in the first twelve weeks from claim established date	X	X			X	X
6	Barriers	80% of claims on the MAVEN FRQ work list receive an FRQ within the first six weeks from claim established date		X				X
7	Opioids	≥ 90% of workers on chronic opioid therapy are dosed at <50mg/day MED		X				
8	Opioids	≥ 90% of worker have an initial opioid prescription of ≤ 3 days (<i>non-surgical comparison groups</i>) OR ≥ 90% of worker have an initial opioid prescription of ≤ 7 days (<i>surgical comparison groups</i>)	X	X	X	X		
9	Opioids	< 5% of workers taking opioids are transitioned to chronic opioid therapy	X	X				
Minimum Best Practices for Medium Adoption			5		3		3	
Minimum Best Practices for High Adoption			9		4		6	